

Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

U. S. DEPT. OF AGRICULTURE
NATIONAL AGRICULTURAL LIBRARY

JUN 23 1967

CURRENT SERIAL RECORDS

1.96
R31Fsm
exp. 2



WATER SUPPLY OUTLOOK FOR COLORADO AND NEW MEXICO

and
FEDERAL - STATE - PRIVATE COOPERATIVE SNOW SURVEYS

UNITED STATES DEPARTMENT of AGRICULTURE...SOIL CONSERVATION SERVICE
and
COLORADO AGRICULTURAL EXPERIMENT STATION
STATE ENGINEER of COLORADO
and STATE ENGINEER of NEW MEXICO

Data included in this report were obtained by the agencies named above in cooperation with the Bureau of Reclamation, U.S. Forest Service, National Park Service, Corps of Engineers and other Federal, State, and private organizations.

AS OF
MAY 1, 1967

TO RECIPIENTS OF WATER SUPPLY OUTLOOK REPORTS:

Most of the usable water in western states originates as mountain snowfall. This snowfall accumulates during the winter and spring, several months before the snow melts and appears as streamflow. Since the runoff from precipitation as snow is delayed, estimates of snowmelt runoff can be made well in advance of its occurrence. Streamflow forecasts published in this report are based principally on measurement of the water equivalent of the mountain snowpack.

Forecasts become more accurate as more of the data affecting runoff are measured. All forecasts assume that climatic factors during the remainder of the snow accumulation and melt season as they affect runoff will add to be an effective average. Early season forecasts are therefore subject to a greater change than those made on later dates.

The snow course measurement is obtained by sampling snow depth and water equivalent at surveyed and marked locations in mountain areas. A total of about ten samples are taken at each location. The average of these are reported as snow depth and water equivalent. These measurements are repeated in the same location near the same dates each year.

Snow surveys are made monthly or semi-monthly from January 1 through June 1 in most states. There are about 1400 snow courses in Western United States and in the Columbia Basin in British Columbia. In the near future, it is anticipated that automatic snow water equivalent sensing devices along with radio telemetry will provide a continuous record of snow water equivalent at key locations.

Detailed data on snow course and soil moisture measurements are presented in state and local reports. Other data on reservoir storage, summaries of precipitation, current streamflow, and soil moisture conditions at valley elevations are also included. The report for Western United States presents a broad picture of water supply outlook conditions, including selected streamflow forecasts, summary of snow accumulation to date, and storage in larger reservoirs.

Snow survey and soil moisture data for the period of record are published by the Soil Conservation Service by states about every five years. Data for the current year is summarized in a West-wide basic data summary and published about October 1 of each year.

Listed below are water supply outlook reports based on Federal-State-Private Cooperative snow surveys. Those published by the Soil Conservation Service may be obtained from Soil Conservation Service, Room 507, Federal Building, 701 N. W. Glisan, Portland, Oregon 97209.

PUBLISHED BY SOIL CONSERVATION SERVICE

D. A. WILLIAMS, Administrator

The Soil Conservation Service publishes reports following the principal snow survey dates from January 1 through June 1 in cooperation with state water administrators, agricultural experiment stations and others. Copies of the reports for Western United States and all state reports may be obtained from Soil Conservation Service, Western Regional Technical Service Center, Room 507, 701 N. W. Glisan, Portland, Oregon 97209.

Copies of state and local reports may also be obtained from state offices of the Soil Conservation Service in the following states:

| STATE | ADDRESS |
|--------------------|---|
| Alaska | P. O. Box "F", Palmer, Alaska 99645 |
| Arizona | 6029 Federal Building, Phoenix, Arizona 85205 |
| Colorado (N. Mex.) | 12417 Federal Building, Denver, Colorado 80202 |
| Idaho | P. O. Box 38, Boise, Idaho 83701 |
| Montana | P. O. Box 855, Bozeman, Montana 59715 |
| Nevada | P. O. Box 4850, Reno Nevada 89505 |
| Oregon | 1218 S. W. Washington St., Portland, Oregon 97205 |
| Utah | 4001 Federal Building, Salt Lake City, Utah 84111 |
| Washington | 840 Bon Marche Bldg., Spokane, Washington 99206 |
| Wyoming | P. O. Box 340, Casper, Wyoming 82602 |

PUBLISHED BY OTHER AGENCIES

Water Supply Outlook reports prepared by other agencies include a report for California by the Water Supply Forecast and Snow Surveys Unit, California Department of Water Resources, P. O. Box 388, Sacramento, California 95802 --- and for British Columbia by the Department of Lands, Forests and Water Resources, Water Resources Service, Parliament Building, Victoria, British Columbia



**FEDERAL-STATE COOPERATIVE
SNOW SURVEYS AND WATER SUPPLY FORECASTS
for
COLORADO RIVER, PLATTE RIVER
ARKANSAS RIVER AND RIO GRANDE DRAINAGE BASINS
issued
May 1, 1967**


Report Prepared By

Jack N. Washichek, Snow Survey Supervisor
and
Donald W. McAndrew, Assistant Snow Survey Supervisor
Fort Collins, Colorado

United States Department of Agriculture
Soil Conservation Service
and
Colorado Agricultural Experiment Station
Fort Collins, Colorado

State Engineer of Colorado
Denver, Colorado
and
State Engineer of New Mexico
Santa Fe, New Mexico

WATER SUPPLY OUTLOOK



THE MAP ON THIS PAGE INDICATES THE MOST PROBABLE WATER SUPPLY AS OF THE DATE OF THIS REPORT. ESTIMATES ASSUME AVERAGE CONDITIONS OF SNOW FALL, PRECIPITATION AND OTHER FACTORS FROM THIS DATE TO THE END OF THE FORECAST PERIOD. AS THE SEASON PROGRESSES ACCURACY OF ESTIMATES IMPROVE. IN ADDITION TO EXPECTED STREAMFLOW, RESERVOIR STORAGE, SOIL MOISTURE IN IRRIGATED AREAS, AND OTHER FACTORS ARE CONSIDERED IN ESTIMATING WATER SUPPLY. ESTIMATES APPLY TO IRRIGATED AREAS ALONG THE MAIN STREAMS AND MAY NOT INDICATE CONDITIONS ON SMALL TRIBUTARIES.



WATER SUPPLY OUTLOOK FOR COLORADO AND NEW MEXICO

as of

May 1, 1967



COLORADO

- All of Colorado's streams are expected to flow less than normal this year, however, the Colorado mainstem and streams to the north should not experience any major shortage. Despite the late spring snow storms, the Colorado snow pack remains below normal.

The South Platte and Arkansas Basins on the east slope will have considerably less than normal streamflow, but have good reservoir storage for supplemental supplies. The Arkansas Basin has 160% of average carryover.

The Rio Grande Basin streams should flow about 65% of normal and storage is below normal. Streams in San Juan Basin should flow 60% of the 15 year average.

Soils are in fair condition throughout most of the state.

Conserve your water!



NEW MEXICO

- Water shortages will exist over most of New Mexico this summer unless spring and summer rainfall is much above normal. Most of the snow in New Mexico is gone. In the headwater areas of the Rio Grande and San Juan Drainages the high elevation snow pack is still good, but medium to low snow has melted.

Reservoir carry-over storage is less than normal, but will be of some help during the summer. Valley soils throughout the state are generally dry.

Forecasts are far less than one-half the normal flows in all streams in New Mexico.

Conservation of water should be the by-word this summer.

TABLE OF CONTENTS

WATER SUPPLY OUTLOOK BY MAJOR WATERSHED AREAS

WATERSHED I

SOUTH PLATTE RIVER WATERSHED

Describes water supply conditions in Fort Collins, Big Thompson, Longmont, Boulder Valley, Jefferson, Teller-Park, Douglas County, Morgan, Kiowa, West Arapahoe, West Adams, East Adams, Platte Valley, Southeast Weld, and West Greeley Soil Conservation Districts.

WATERSHED II

ARKANSAS RIVER WATERSHED

Describes water supply conditions in Lake County, Upper Arkansas, Fremont, Custer County Divide, Fountain Valley, Black Squirrel, Horse-Rush Creek, Central Colorado, Turkey Creek, Pueblo, Bessemer, Olney Boone, Cheyenne, Upper Huerfano, Stonewall, Spanish Peaks, Purgatoire, Branson Trinchera, Western Baca County, Southeastern Baca County, Two Buttes, Bent, Timpas, Northeast Prowers, Prowers, West Otero, East Otero, and Big Sandy Soil Conservation Districts.

WATERSHED III

RIO GRANDE WATERSHED (COLORADO)

Describes water supply conditions in Rio Grande, Center, Mosca Hooper, Mt. Blanca, Sanches, and Culebra Soil Conservation Districts.

WATERSHED IV

RIO GRANDE WATERSHED (NEW MEXICO)

Describes water supply conditions in Lower Cebolla, Abiquiu-Vallecitos, Eastern Taos, Lindrieth, Coyote-Canones, Espanola Valley, Pojoaque, Jemez, Santa Fe-Sandoval, Tijeras, Cuba, and Englewood Soil Conservation Districts.

WATERSHED V

DOLORES, SAN JUAN, AND ANIMAS RIVERS WATERSHED

Describes water supply conditions in San Miguel Basin, Dove Creek, Dolores, Mancos, LaPlata, Pine River, San Juan, and Glade Park Soil Conservation Districts.

WATERSHED VI

GUNNISON RIVER WATERSHED

Describes water supply conditions in Delta, Gunnison, Cimarron, Shavano, and Uncompahgre Soil Conservation Districts.

WATERSHED VII

COLORADO RIVER WATERSHED

Describes water supply conditions in DeBeque, Lower Grand Valley, Bookcliff, Eagle County, Middle Park, Glade Park, Upper Grand Valley, Plateau Valley, South Side, and Mt. Sopris Soil Conservation Districts.

WATERSHED VIII

YAMPA, WHITE AND NORTH PLATTE RIVERS WATERSHED

Describes water supply conditions in Yampa, Moffat, West Routt, East Routt, North Park, Upper White River, Lower White River, and Douglas Creek Soil Conservation Districts.

WATERSHED IX


LOWER SOUTH PLATTE RIVER WATERSHED

Describes water supply conditions in Sedgwick, South Platte, Haxton Peetz, Padroni, Morgan, Rock Creek and Yuma Soil Conservation Districts.

WATERSHED I

WATER SUPPLY OUTLOOK
FOR THE SOIL CONSERVATION DISTRICTS IN THE
SOUTH PLATTE RIVER WATERSHED IN COLORADO
as of
May 1, 1967

U. S. DEPARTMENT OF AGRICULTURE - SOIL CONSERVATION SERVICE
COLORADO EXPERIMENT STATION - STATE ENGINEERS OF COLORADO AND NEW MEXICO



The snow pack in the mountain watersheds of the South Platte River and its tributaries remains below normal. Even with the big storm of April 13th, the snow pack remains at only 67% of average over the entire basin. There are a few isolated areas in the basin that have a near normal snow pack. These are mostly in the very high elevations.

The water held in the reservoirs throughout the basin remains similar to last month at 107% of average. This water will be an excellent supplement this summer. Following the good rains in the area during this last month most of the irrigated areas are reporting good soil moisture conditions. This situation will help the below normal streamflow as it will lessen the demands on the early flows.

Mountain soil moisture is slightly below normal for this time of year. Some of the snow water will be used to wet up the soil mantle before the spring runoff starts.

Streamflow forecasts range from a high of 90% of average on Clear Creek to a low of 73% on the Cache La Poudre. The Big Thompson, Boulder Creek and Saint Vrain Rivers will flow between 77 to 85% this year. The mainstem of the South Platte will probably flow less than 70% this summer.

Issued By: Soil Conservation Service

F. A. Mark, State Conservationist,
Colorado

E. A. Nicholson, Area Conservationist,
Littleton, Colorado

"THE CONSERVATION OF WATER BEGINS WITH THE SNOW SURVEY"

SNOW

CURRENT INFORMATION

PAST RECORD

| Snow Course | Date of Survey | Snow Depth (Inches) | Water Content (Inches) | Water Content (Inches) | |
|----------------------------------|----------------------|---------------------------|------------------------------|---------------------------|---------------|
| | | | | Last Year | Avg. 48-62 |
| South Platte River & Tributaries | | | | | |
| Baltimore | 4/27 | 0 | 0 | 0 | - - |
| Berthoud Falls | 4/27 | 19 | 7.2 | 8.4 | 13.8* |
| Big South | 4/29 | 3 | 0.4 | 0 | 0.8 |
| Boulder Falls | 4/29 | 22 | 9.2 | 5.0 | 13.2* |
| Cameron Pass | 4/26 | 82 | 33.9 | 21.9 | 28.1 |
| Chambers Lake | 4/30 | 14 | 6.1 | 0 | 5.5 |
| Como | 4/26 | 6 | 2.1 | | - - |
| Copeland Lake | 4/28 | 0 | 0.0 | 0 | 2.3* |
| Deadman Hill | 4/27 | 48 | 17.1 | 13.0 | 18.1 |
| Deer Ridge | 4/27 | 1 | 0.5 | 0 | 3.5* |
| Empire | 4/27 | 22 | 8.4 | 4.2 | 7.1* |
| Geneva Park | 4/26 | 5 | 2.1 | 0.5 | 1.9* |
| Grizzly Peak (B) | 4/27 | 49 | 18.4 | 10.7 | 21.1 |
| Hidden Valley | 4/27 | 29 | 10.6 | 6.9 | 13.6 |
| Hoosier Pass | 4/26 | 33 | 11.6 | 6.0 | 12.9 |
| Horseshoe | 4/26 | 18 | 6.0 | | - - |
| Hour Glass Lake | 4/27 | 12 | 4.1 | 1.4 | 7.5 |
| Jefferson Creek | 4/27 | 17 | 5.8 | 1.9 | 8.0* |
| Lake Irene (B) | 4/26 | 61 | 21.8 | 12.0 | 24.7 |
| Long's Peak | 4/29 | 33 | 12.1 | 6.5 | 13.4* |
| Lost Lake | 4/30 | 32 | 7.7 | 1.9 | 10.2* |
| Loveland Lift No. 1 | 4/28 | 76 | 27.5 | 15.1 | - - |
| Loveland Pass | 4/28 | 33 | 13.7 | 4.9 | 16.4 |
| Mosquito | 4/26 | 3 | 1.1 | | - - |
| Pine Creek | 4/27 | 0 | 0.0 | 0 | - - |
| Red Feather | | | | 0.7 | 4.9* |
| Two Mile | 4/27 | 53 | 17.3 | 9.8 | 17.8* |
| Trout Creek | 4/28 | 0 | 0.0 | | - - |
| University Camp | 4/29 | 35 | 14.1 | 8.2 | 24.9 |
| Ward | 4/27 | 11 | 4.1 | 0.4 | 6.0* |
| Wild Basin | 4/27 | 24 | 7.6 | 5.1 | 14.8 |

SOIL MOISTURE

| Station | Date of Survey | Capacity (Inches) | This Year | Last Year | Avg. All Data |
|---------------|----------------|-------------------|-----------|-----------|---------------|
| Alpine Camp | 4/27 | 6.9 | 3.5 | 4.1 | 4.3 |
| Beaver Dam | 4/27 | 7.3 | 4.8 | 5.2 | 4.7 |
| Clear Creek | 4/28 | 9.5 | 5.8 | 6.4 | 5.9 |
| Feather | 4/28 | 10.1 | 6.5 | 9.4 | 8.1 |
| Guard Station | 4/29 | 6.9 | 4.9 | 4.6 | 4.7 |
| Hoop Creek | 4/26 | 4.9 | 3.5 | 3.5 | 2.9 |
| Hoosier Pass | 4/26 | 7.8 | 4.8 | 6.3 | 5.9 |
| Kenosha Pass | 4/27 | 4.4 | 4.0 | 3.3 | 3.7 |
| Laramie Road | 4/30 | 12.4 | 8.1 | 9.1 | 9.0 |
| Two Mile | 4/27 | 9.1 | 4.3 | 5.5 | 5.6 |

ALL PROFILES 4 FEET DEEP

RETURN IF NOT DELIVERED

UNITED STATES

DEPARTMENT OF AGRICULTURE

SOIL CONSERVATION SERVICE

Snow Survey
Colorado State University
Fort Collins, Colorado

OFFICIAL BUSINESS

RESERVOIR STORAGE (1,000 Acre-Feet)

| Reservoir | Usable Capacity | This Year | Last Year | 15 Year Average 1948-62 |
|-----------------|-----------------|-----------|-----------|-------------------------|
| Antero | 33.0 | 14.5 | 15.9 | 13.4 |
| Barr Lake | 32.2 | 15.7 | 28.3 | 24.7 |
| Black Hollow | 8.0 | 3.3 | 4.1 | 3.3 |
| Boyd Lake | 58.0 | 28.5 | 40.6 | 20.8 |
| Cache La Poudre | 9.5 | 8.3 | 9.0 | 7.7 |
| Carter Lake | 108.9 | 95.7 | 107.3 | 79.0 |
| Chambers Lake | 8.8 | 3.2 | 6.7 | 2.8 |
| Cheeseman | 79.0 | 31.8 | 77.2 | 54.3 |
| Cobb Lake | 34.3 | 0 | 7.3 | 9.2 |
| Eleven Mile | 81.9 | 90.9 | 92.3 | 74.6 |
| Fossil Creek | 11.6 | 8.0 | 10.1 | 7.1 |
| Gross | 43.1 | 19.0 | 24.2 | - - |
| Halligan | 6.4 | 6.4 | 6.4 | 3.9 |
| Horsetooth | 143.5 | 116.8 | 120.3 | 85.6 |
| Lake Loveland | 13.6 | 4.1 | 8.3 | 7.4 |
| Lone Tree | 9.2 | 5.4 | 8.3 | 7.9 |
| Mariano | 5.4 | 5.3 | 5.6 | 3.2 |
| Marshall | 10.3 | 2.1 | 7.4 | 4.4 |
| Marston | 18.9 | 15.9 | 15.3 | 15.2 |
| Milton | 24.4 | 6.9 | 18.4 | 12.5 |
| Standly | 18.5 | 9.8 | 20.6 | 12.6 |
| Terry Lake | 8.2 | 4.3 | 5.9 | 5.2 |
| Union | 12.7 | 6.8 | 12.7 | 8.2 |
| Windsor | 18.6 | 6.3 | 13.1 | 11.4 |

MEASURED FIRST OF MONTH

STREAMFLOW FORECAST (1,000 Acre-Feet)

| Stream and Station | Forecast Period April - Sept. | This Year % of Avg. | Avg. 1948-1962 |
|------------------------------------|-------------------------------|---------------------|----------------|
| Big Thompson at Drake (2) | 85 | 77 | 110 |
| Boulder at Orde11 | 46 | 85 | 54 |
| Cache La Poudre at Canon Mouth (1) | 180 | 73 | 246 |
| Clear Creek at Golden (3) | 120 | 90 | 134 |
| Saint Vrain at Lyons | 65 | 81 | 80 |

- (1) Observed flow minus diversions from Michigan, Colorado and Laramie Rivers, plus diversions for irrigation and municipal use above station.
 (2) Observed flow plus by-pass to power plants.
 (3) Observed flow minus diversions through Jones Pass.

NOTE: * - 1948-62 (adjusted average)

NS - NO SURVEY

(A) - AIR OBSERVED

(B) - ON ADJACENT DRAINAGE


This Report Prepared by Jack N. Washichek and Donald W. McAndrew, Soil Conservation Service, Colorado State University, Fort Collins, Colo.

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF AGRICULTURE

WATERSHED II

WATER SUPPLY OUTLOOK
FOR THE SOIL CONSERVATION DISTRICTS IN THE
ARKANSAS RIVER WATERSHED IN COLORADO
as of
May 1, 1967

U. S. DEPARTMENT OF AGRICULTURE - SOIL CONSERVATION SERVICE
COLORADO EXPERIMENT STATION - STATE ENGINEERS OF COLORADO AND NEW MEXICO



The mountain snow pack on the Arkansas River Drainage is only 57% of average. However, the snow ranges from exactly normal in the Fremont Pass area to much below normal in the southern mountains. The snow courses located in the headwaters area of the Purgatoire and Cucharas Drainages are completely void of snow.

The soil moisture conditions in the mountains is better than average and similar to last year. Low elevation snow melt probably accounts for this condition. Soil moisture in the irrigated areas however, is reported to be only poor to fair. As of this writing the streamflow is below normal in the Arkansas River and its tributaries.

Water held in storage is still the only bright spot in the water outlook for the Arkansas Valley. Currently storage is 160% of average. There is nearly 300,000 acre-feet of water stored in the major reservoirs in the Arkansas River Watershed, however, nearly 140,000 acre-feet of this is in John Martin. This reservoir storage will be an excellent supplement for the deficient streamflow this season.

Streamflow forecasts range from a low of 38% of average on the Purgatoire to 65% of normal for the Arkansas at Salida. The streams in the wet mountain valley area will flow only half of normal this year. Streamflow on the Huerfano, Cucharas and Purgatoire Rivers will be about 40% of average this year.

Issued By: Soil Conservation Service

F. A. Mark, State Conservationist,
Colorado

Will D. McCorkle, Area Conservationist,
La Junta, Colorado

"THE CONSERVATION OF WATER BEGINS WITH THE SNOW SURVEY"

SNOW

CURRENT INFORMATION

PAST RECORD

| Snow Course | Date of Survey | Snow Depth (Inches) | Water Content (Inches) | Water Content (Inches) | |
|-------------------|----------------|---------------------|------------------------|------------------------|------------|
| | | | | Last Year | Avg. 48-62 |
| Arkansas River | | | | | |
| Bigelow Divide | 4/28 | 0 | 0 | 0 | - - |
| Bourbon | 4/27 | 0 | 0 | 0 | 2.9* |
| Cooper Hill | 4/26 | 43 | 12.6 | 6.8 | - - |
| Cucharas Pass | 4/28 | 0 | 0 | 1.6 | - - |
| East Fork | 4/26 | 16 | 5.2 | 2.0 | 13.4 |
| Four Mile Park | 4/28 | 0 | 0 | 0 | 1.0 |
| Fremont Pass | 4/26 | 57 | 19.5 | 10.3 | 19.5 |
| Garfield | 4/28 | 3 | 1.5 | 3.1 | - - |
| LaVeta Pass (B) | 4/28 | 0 | 0 | 0 | 1.7 |
| Monarch Pass | 4/28 | 14 | 6.8 | 9.3 | 18.4 |
| St. Elmo (Air) | 4/27 | 19 | 6.7 | 5.6 | 11.8 |
| Tennessee Pass | 4/28 | 8 | 2.7 | 5.0 | 8.5 |
| Tomichi | 4/28 | 6 | 2.2 | 5.5 | - - |
| Twin Lakes Tunnel | 4/28 | 24 | 8.8 | 4.0 | 9.1 |
| Westcliffe | 4/28 | 0 | 0 | 0 | 1.1 |

SOIL MOISTURE

| Station | Date of Survey | Capacity (Inches) | This Year | Last Year | Avg. All Data |
|-------------------|----------------|-------------------|-----------|-----------|---------------|
| Garfield | 4/26 | 6.7 | 6.1 | 6.3 | 4.3 |
| King | 4/28 | 3.3 | 3.0 | 3.0 | 2.1 |
| LaVeta Pass | 4/28 | 11.9 | 11.7 | 11.9 | 11.8 |
| Leadville | 4/26 | 7.8 | 5.7 | 5.7 | 4.8 |
| Twin Lakes Tunnel | 4/26 | 4.5 | 2.9 | 4.2 | 3.1 |

ALL PROFILES 4 FEET DEEP

RESERVOIR STORAGE (1,000 Acre-Feet)

| Reservoir | Usable Capacity | This Year | Last Year | 15 Year Average 1948-62 |
|--------------|-----------------|-----------|-----------|-------------------------|
| Adobe Creek | 61.6 | 21.5 | 54.4 | 13.0 |
| Clear Creek | 11.4 | 6.8 | 11.2 | 4.7 |
| Cucharas | 40.0 | 0.1 | 0 | 5.3 |
| Great Plains | 150.0 | 71.8 | 123.3 | 44.4 |
| Horse Creek | 26.9 | 5.7 | 21.0 | 5.6 |
| John Martin | 366.6 | 136.9 | 352.5 | 64.6 |
| Meredith | 41.9 | 4.6 | 24.5 | 10.4 |
| Model | 15.0 | - | - | 2.2 |
| Sugar Loaf | 17.4 | 8.7 | 11.3 | 6.8 |
| Twin Lakes | 57.9 | 18.3 | 41.0 | 17.2 |

MEASURED FIRST OF MONTH

STREAMFLOW FORECAST (1,000 Acre-Feet)

| Stream and Station | Forecast Period April - Sept. | This Year % of Avg. | Avg. 1948-1962 |
|------------------------|-------------------------------|---------------------|----------------|
| Arkansas at Pueblo (4) | 194 | 60 | 323 |
| Arkansas at Salida (4) | 225 | 65 | 345 |
| Cucharas nr LaVeta | 6 | 43 | 14 |
| Purgatoire at Trinidad | 17 | 38 | 45 |

(4) Observed flow plus change in Clear Creek, Twin Lakes, and Sugar Loaf Reservoirs minus diversions through Busk-Ivanhoe and Twin Lake Tunnels and Ewing, Fremont Pass, Wurtz and Columbine Ditches.

NOTE: * - 1948-62 (adjusted averages)

NS - NO SURVEY

(A) - AIR OBSERVED

(B) - ON ADJACENT DRAINAGE

This Report Prepared by Jack N. Washichek and Donald W. McAndrew, Soil Conservation Service, Colorado State University, Fort Collins, Colo.

RETURN IF NOT DELIVERED

UNITED STATES

DEPARTMENT OF AGRICULTURE

SOIL CONSERVATION SERVICE

Snow Survey
Colorado State University
Fort Collins, Colorado


OFFICIAL BUSINESS

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF AGRICULTURE

WATERSHED III

WATER SUPPLY OUTLOOK
FOR THE SOIL CONSERVATION DISTRICTS IN THE
UPPER RIO GRANDE WATERSHED IN COLORADO
as of
May 1, 1967

U. S. DEPARTMENT OF AGRICULTURE - SOIL CONSERVATION SERVICE
COLORADO EXPERIMENT STATION - STATE ENGINEERS OF COLORADO AND NEW MEXICO



Streamflow will be less than normal and unless spring and summer rainfall is above normal, water shortages will exist. Flows should drop off rapidly and late season streamflow will be very low.

All the low to medium elevation snow pack is gone. The snow line now is about 10,000 feet. The snow pack above 10,000 feet is nearly normal. This high elevation snow pack will not provide adequate runoff this summer.

Practically all of the snow pack in the Sangre De Cristo Range is gone and still streamflow remains near normal.

Carry-over storage in the major reservoirs in the basin contain 74% of normal storage.

Mountain soils are drier than normal despite the snow melt. This will tend to reduce runoff still further.

Forecasts are computed assuming normal precipitation for the remainder of the year.

The Rio Grande in Colorado should flow near 65% of the 1948-62 average. The Alamosa and South Fork should flow about 70%. Streams originating in the Sangre De Cristo Range will flow far below normal.

Water will have to be conserved to the utmost this summer.

Issued By: Soil Conservation Service

F. A. Mark, State Conservationist,
Colorado

R. K. Griffin, Area Conservationist,
Durango, Colorado

"THE CONSERVATION OF WATER BEGINS WITH THE SNOW SURVEY"

SNOW

CURRENT INFORMATION

PAST RECORD

| Snow Course | Date of Survey | Snow Depth (Inches) | Water Content (Inches) | Water Content (Inches) | |
|-------------------------------|----------------|---------------------|------------------------|------------------------|------------|
| | | | | Last Year | Avg. 48-62 |
| <u>Rio Grande in Colorado</u> | | | | | |
| Cochetopa Pass | 4/27 | 0 | 0 | 0 | 2.7* |
| Hiway | 4/27 | 64 | 25.4 | 25.7 | 27.8* |
| Lake Humphreys | 4/26 | 0 | 0 | 0.4 | 0.2* |
| Pass Creek | 4/27 | 0 | 0 | 2.2 | 3.3* |
| Pool Table | 4/26 | 0 | 0 | 1.8 | 1.9* |
| Porcupine | 4/25 | 8 | 2.5 | 7.4 | 6.8* |
| Red Mountain Pass (B) | 4/25 | 61 | 25.0 | 25.5 | 31.4* |
| Santa Maria | 4/28 | 0 | 0 | 0 | 0.5 |
| Upper Rio Grande | 4/27 | 0 | 0 | 1.8 | 2.3 |
| Wolf Creek Pass | 4/27 | 52 | 24.5 | 23.2 | 24.7 |
| Wolf Creek Summit (B) | 4/27 | 73 | 29.3 | 32.2 | 30.2 |
| <u>Alamosa River</u> | | | | | |
| Silver Lakes | 4/27 | 0 | 0 | 0 | 0.5 |
| Summitville | 4/24 | 54 | 17.4 | 17.6 | 20.5 |
| <u>Conejos River</u> | | | | | |
| Cumbres Pass | 4/29 | 29 | 14.4 | 11.8 | 12.5 |
| Platoro | 4/27 | 30 | 11.5 | 9.2 | 10.9* |
| River Springs | 4/27 | 0 | 0 | - | 0.7 |
| <u>Sangre De Cristo Range</u> | | | | | |
| Cucharas Pass (B) | 4/28 | 0 | 0 | 1.6 | - |
| Culebra | 4/28 | 1 | 0.4 | 5.3 | 5.2 |
| LaVeta Pass | 4/28 | 0 | 0 | 0 | 1.7 |

SOIL MOISTURE

| Station | Date of Survey | Capacity (Inches) | This Year | | Avg. All Data |
|--------------|----------------|-------------------|-----------|------|---------------|
| | | | Year | Year | |
| Alberta Park | 4/24 | 8.2 | 5.7 | 7.1 | 5.6 |
| Bristol View | 4/26 | 6.1 | 3.7 | 6.1 | 4.4 |
| LaVeta Pass | 4/28 | 11.9 | 11.7 | 11.9 | 11.8 |
| Mogote | 4/27 | 10.7 | 8.2 | 8.6 | 9.0 |

ALL PROFILES 4 FEET DEEP

RESERVOIR STORAGE (1,000 Acre-Feet)

| Reservoir | Usable Capacity | This Year | Last Year | 15 Year Average 1948-62 |
|-------------|-----------------|-----------|-----------|-------------------------|
| Continental | 26.7 | 5.2 | 10.1 | 7.7 |
| Platoro | 60.0 | 3.0 | 17.3 | - |
| Rio Grande | 45.8 | 10.3 | 39.8 | 14.8 |
| Sanchez | 103.2 | 9.9 | 15.1 | 12.3 |
| Santa Maria | 45.0 | 3.6 | 18.4 | 7.8 |
| Terrace | 17.7 | 5.9 | 10.8 | 4.8 |

MEASURED FIRST OF MONTH

STREAMFLOW FORECAST (1,000 Acre-Feet)

| Stream and Station | Forecast Period | This Year | Avg. |
|----------------------------------|-----------------|-----------|-----------|
| | April - Sept. | % of Avg. | 1948-1962 |
| Alamosa abv Terrace | 48 | 71 | 68 |
| Conejos nr Mogote | 130 | 66 | 196 |
| Culebra at San Luis (6) | 8 | 38 | 21 |
| Rio Grande at 30 Mile Bridge (5) | 85 | 64 | 132 |
| Rio Grande nr Del Norte (5) | 275 | 56 | 492 |
| South Fork at South Fork | 80 | 66 | 122 |

- (5) Observed flow plus change in storage in Santa Maria, Rio Grande and Continental Reservoir.
 (6) Observed flow plus changes in storage in Sanchez Reservoir.

NOTE: * - 1948-62 (adjusted averages)
 NS - NO SURVEY
 (A) - AIR OBSERVED
 (B) - ON ADJACENT DRAINAGE

This Report Prepared by Jack N. Washichek and Donald W. McAndrew, Soil Conservation Service, Colorado State University, Fort Collins, Colo.

RETURN IF NOT DELIVERED

UNITED STATES

DEPARTMENT OF AGRICULTURE

SOIL CONSERVATION SERVICE

Snow Survey
 Colorado State University
 Fort Collins, Colorado


OFFICIAL BUSINESS

POSTAGE AND FEES PAID
 U.S. DEPARTMENT OF AGRICULTURE

RIO GRANDE WATERSHED IN NEW MEXICO

as of
May 1, 1967

U. S. DEPARTMENT OF AGRICULTURE - SOIL CONSERVATION SERVICE
COLORADO EXPERIMENT STATION - STATE ENGINEERS OF COLORADO AND NEW MEXICO



April snowfall did not improve the water supply outlook in New Mexico. All but the extremely high snow pack is gone in New Mexico. The snow in the headwaters of the Rio Grande in Colorado is also melted below 10,000 feet. Although the extremely high elevation snow pack is good it will not hold up summer flows.

Carry-over storage in major reservoirs on the Rio Grande is 65% of normal, but will still be a good supplement.

Storage in Conchas is 164,500 acre-feet compared to a normal of 229,500 acre-feet.

The two reservoirs on the Pecos River, Alamogordo and McMillan-Avalon, contain 74,300 acre-feet which is almost exactly normal for this date.

Mountain soils are wetter than normal, which may be caused by the early snow melt.

Valley soils in the upper basin contain only fair amounts of moisture, while all other areas of the State report dry soils.

Forecasts of summer streamflow are all less than half of normal.

Strict conservation measures will have to be employed by water users throughout the State this summer.

Issued By: Soil Conservation Service

Einar L. Roget, State Conservationist,
Albuquerque, New Mexico

Walter B. Rumsey, Area Conservationist,
Santa Fe, New Mexico

"THE CONSERVATION OF WATER BEGINS WITH THE SNOW SURVEY"

| SNOW | CURRENT INFORMATION | | | PAST RECORD | |
|--------------------------|---------------------|---------------------|------------------------|------------------------|------------|
| Snow Course | Date of Survey | Snow Depth (Inches) | Water Content (Inches) | Water Content (Inches) | |
| | | | | Last Year | Avg. 48-62 |
| Rio Grande (Colorado) | | | | | |
| Culebra | 4/28 | 1 | 0.4 | 5.3 | 5.2 |
| Cumbres Pass | 4/29 | 29 | 14.4 | 11.8 | 12.5 |
| LaVeta Pass | 4/28 | 0 | 0.0 | 0 | 1.7 |
| Platoro | 4/27 | 30 | 11.5 | 9.2 | 10.9* |
| River Springs | 4/27 | 0 | 0.0 | | 0.7 |
| Santa Maria | 4/28 | 0 | 0.0 | 0 | 0.5 |
| Silver Lakes | 4/27 | 0 | 0.0 | 0 | 0.5 |
| Summitville | 4/24 | 54 | 17.4 | 17.6 | 20.5 |
| Upper Rio Grande | 4/27 | 0 | 0.0 | 1.8 | 2.3 |
| Wolf Creek Pass | 4/27 | 52 | 24.5 | - - | - - |
| Big Tesuque (New Mexico) | | | | | |
| Chamita | 4/27 | 0 | 0.0 | - - | - - |

RESERVOIR STORAGE (1,000 Acre-Feet)

| Reservoir | Usable Capacity | This Year | Last Year | 15 Year Average 1948-62 |
|------------------|-----------------|-----------|-----------|-------------------------|
| Alamogordo | 122.1 | 69.0 | 27.8 | 63.8 |
| Caballo | 344.0 | 94.3 | 103.4 | 102.1 |
| Conchas | 280.3 | 164.5 | 239.3 | 229.5 |
| Elephant Butte | 2206.8 | 222.8 | 479.4 | 354.0 |
| El Vado | 194.5 | 13.6 | 13.0 | 55.1 |
| McMillan-Avalon | 37.0 | 5.3 | 11.5 | 10.6 |
| Red Bluff (Tex.) | 307.0 | | 52.7 | 59.1 |

MEASURED FIRST OF MONTH

STREAMFLOW FORECAST(1,000 Acre-Feet)

| Stream and Station | Forecast as Indicated | This Year % of Avg. | Avg. 1948 - 62 |
|--------------------------------|-----------------------|---------------------|----------------|
| Costilla at Costilla (8) | 8 AS | 32 | 25 |
| Pecos at Pecos | 20 AS | 38 | 53 |
| Rio Chama nr La Puente | 100 AS | 47 | 214 |
| Rio Grande at Otowi (7)* | 250 MJ | 41 | 609 |
| Rio Grande at San Marcial (7)* | 95 MJ | 22 | 424 |
| Rio Hondo nr Valdez | 9 AS | 50 | 18 |
| Red River at Questa** | 9 AJ | 36 | 25 |

The Forecast of the Rio Grande at San Marcial is 14 % of the Average used by the Elephant Butte Irrigation District.

A-S is April through September.

A-J is April through July.

M-J is March through July.

- (7) Observed flow plus changes in storage in El Vado and Abiquiu Reservoirs.
 (8) Observed flow plus changes in storage in Costilla Reservoir.

NOTE: * - 1948-62 (adjusted averages)

NS - NO SURVEY

(A) - AIR OBSERVED

(B) - ON ADJACENT DRAINAGE

This Report Prepared by Jack N. Washichek and Donald W. McAndrew, Soil Conservation Service, Colorado State University, Fort Collins, Colo.

SOIL MOISTURE

| Station | Date of Survey | Capacity (Inches) | This Year | Last Year | Avg. All Data |
|-------------------|----------------|-------------------|-----------|-----------|---------------|
| Colorado | | | | | |
| Alberta Park | 4/24 | 8.2 | 5.7 | 7.1 | 5.6 |
| Bristol View | 4/26 | 6.1 | 3.7 | 6.1 | 4.4 |
| Mogote | 4/27 | 10.7 | 8.2 | 8.6 | 9.0 |
| New Mexico | | | | | |
| Aqua Piedra | 3/29 | 7.2 | 5.1 | 5.3 | 3.7 |
| Bateman | 3/22 | 6.7 | 4.5 | 4.8 | 2.6 |
| Big Tesuque | 3/31 | 3.7 | 3.3 | 1.9 | 1.7 |
| Chamita | 3/30 | 8.0 | 8.0 | 8.0 | 3.7 |
| Fenton Hill | | 6.5 | - - | 6.5 | 4.5 |
| Red Summit | 3/28 | 4.8 | 1.5 | 1.5 | 2.1 |
| Rio En Medio | 3/31 | 3.5 | 1.0 | 1.6 | 1.1 |
| Taos Canyon | 3/29 | 3.3 | 2.5 | 2.5 | 2.3 |

RETURN IF NOT DELIVERED
UNITED STATES

ALL PROFILES 4 FEET DEEP

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF AGRICULTURE

DEPARTMENT OF AGRICULTURE

SOIL CONSERVATION SERVICE


Snow Survey
Colorado State University
Fort Collins, Colorado

OFFICIAL BUSINESS

WATER SUPPLY OUTLOOK
FOR THE SOIL CONSERVATION DISTRICTS IN THE
SAN MIGUEL - DOLORES - ANIMAS - SAN JUAN
WATERSHEDS IN COLORADO AND NEW MEXICO

as of
May 1, 1967

U. S. DEPARTMENT OF AGRICULTURE - SOIL CONSERVATION SERVICE
COLORADO EXPERIMENT STATION - STATE ENGINEERS OF COLORADO AND NEW MEXICO



High elevation snow pack remains good, but low snow has already melted. Snow pack on the San Juan stands at 87% of normal, however, this is not a true picture of conditions. Only high elevation snows are used in this average. The snow pack on the Animas River is 43% of normal and on the Dolores only 30%.

Low and medium elevation snow pack has already melted. Streamflow has not increased materially with all the low snows gone.

Carry-over reservoir storage is about normal in Groundhog and Vallecito Reservoirs, Navajo Reservoir contains 379,000 acre-feet as of May 1. This will be an excellent supplement for water users under that irrigation system.

Mountain soils contain slightly more moisture than usual. This may be caused by the early snow melt. Some shortages will exist in this area unless spring and summer rains are above normal. The San Juan, Animas, and Dolores Rivers are expected to flow only about 60% of normal. Piedra Creek and Los Pinos should flow slightly better, possibly as high as 70% of the 1948-62 average.

Forecasts are based on normal precipitation during the remainder of the forecast period.

The conservation of water is a must in this area this summer.

Issued By: Soil Conservation Service

F. A. Mark, State Conservationist,
Colorado

E. L. Roget, State Conservationist,
Albuquerque, New Mexico

R. K. Griffin, Area Conservationist,
Durango, Colorado

W. B. Rumsey, Area Conservationist,
Santa Fe, New Mexico

D. B. Beach, Area Conservationist,
Grand Junction, Colorado

"THE CONSERVATION OF WATER BEGINS WITH THE SNOW SURVEY"

SNOW

CURRENT INFORMATION

PAST RECORD

| Snow Course | Date of Survey | Snow Depth (Inches) | Water Content (Inches) | Water Content (Inches) | |
|------------------------|----------------|---------------------|------------------------|------------------------|------------|
| | | | | Last Year | Avg. 48-62 |
| <u>San Juan River</u> | | | | | |
| Chama Divide (B) | 4/28 | 0 | 0 | 0 | - - |
| Chamita (B) | 4/28 | 0 | 0 | 0 | - - |
| Upper San Juan | 4/27 | 48 | 21.0 | 23.1 | 30.2 |
| Wolf Creek Pass (B) | 4/27 | 52 | 24.5 | 23.2 | 24.7 |
| Wolf Creek Summit | 4/27 | 73 | 29.3 | 32.2 | 30.2 |
| <u>Animas River</u> | | | | | |
| Cascade | 4/28 | 0 | 0 | 2.8 | 3.0 |
| Howardville | Destroyed | | | 9.0 | 7.4* |
| Iron-ton Park (B) | 4/27 | 0 | 0 | 0 | 7.1 |
| Mineral Creek | 4/28 | 0 | 0 | 11.3 | 12.1* |
| Molas Lake | 4/28 | 0 | 0 | 5.4 | 7.8* |
| Red Mountain Pass | 4/28 | 61 | 25.0 | 25.5 | 31.4* |
| Silver-ton Sub-Station | 4/28 | 0 | 0 | 0 | 0.1 |
| Spud Mountain | 4/28 | 36 | 14.6 | 18.2 | 23.8* |
| <u>Dolores River</u> | | | | | |
| Lizzard Head | 4/28 | 15 | 6.3 | 13.2 | 13.7 |
| Rico | 4/28 | 0 | 0 | 0 | 1.0 |
| Telluride | 4/27 | 0 | 0 | 0 | 0.7 |
| Trout Lake | 4/27 | 3 | 1.3 | 3.6 | 9.9* |

RESERVOIR STORAGE (1,000 Acre-Feet)

| Reservoir | Usable Capacity | This Year | Last Year | 15 Year Average 1948-62 |
|-----------|-----------------|-----------|-----------|-------------------------|
| Groundhog | 21.7 | 10.3 | 21.7 | 8.6 |
| Navajo | 1036.0 | 379.0 | 255.5 | - - |
| Vallecito | 126.3 | 56.0 | 88.0 | 50.9 |

MEASURED FIRST OF MONTH

STREAMFLOW FORECAST (1,000 Acre-Feet)

| Stream and Station | Forecast Period | This Year | Avg. |
|---------------------------|-----------------|-----------|-----------|
| | April - Sept. | % of Avg. | 1948-1962 |
| Animas at Durango | 280 | 61 | 456 |
| Dolores at Dolores | 150 | 58 | 260 |
| La Plata at Hesperus | 17 | 63 | 27 |
| Los Pinos at Bayfield (9) | 155 | 70 | 220 |
| Piedra Creek nr Piedra | 120 | 66 | 182 |
| San Juan at Rosa (9) | 360 | 60 | 597 |

SOIL MOISTURE

| Station | Date of Survey | Capacity (Inches) | This Year | Last Year | Avg. All Data |
|---------------|----------------|-------------------|-----------|-----------|---------------|
| Cascade | 4/28 | 9.1 | 8.6 | 9.1 | 6.8 |
| Dolores | 4/28 | 19.6 | 12.7 | 15.1 | 11.4 |
| Lizzard Head | 4/28 | 11.8 | 7.6 | 8.6 | 8.5 |
| Mineral Creek | 4/28 | 5.7 | 5.4 | 5.6 | 4.1 |
| Molas Lake | 4/28 | 9.4 | 6.4 | 9.4 | 5.8 |
| Rico | 4/28 | 13.8 | 9.7 | 13.8 | 9.0 |

ALL PROFILES 4 FEET DEEP

(9) Observed flow plus changes in storage in Vallecito Reservoir.

NOTE: * - 1948-62 (adjusted averages)
 NS - NO SURVEY
 (A) - AIR OBSERVED
 (B) - ON ADJACENT DRAINAGE

This Report Prepared by Jack N. Washichek and Donald W. McAndrew, Soil Conservation Service, Colorado State University, Fort Collins, Colo.


RETURN IF NOT DELIVERED
 UNITED STATES
 DEPARTMENT OF AGRICULTURE
 SOIL CONSERVATION SERVICE
 Snow Survey
 Colorado State University
 Fort Collins, Colorado
 OFFICIAL BUSINESS

POSTAGE AND FEES PAID
 U.S. DEPARTMENT OF AGRICULTURE

WATER SUPPLY OUTLOOK
FOR THE SOIL CONSERVATION DISTRICTS IN THE
GUNNISON RIVER WATERSHED IN COLORADO
as of
May 1, 1967

WATERSHED VI

U. S. DEPARTMENT OF AGRICULTURE - SOIL CONSERVATION SERVICE
COLORADO EXPERIMENT STATION - STATE ENGINEERS OF COLORADO AND NEW MEXICO



Again this month the snow pack decreased throughout the headwaters of the Gunnison River. Snow courses in the Lake City, Monarch Pass, and Taylor Park areas are currently about 50% of normal. In this area there has been less snow than now, once in recent history. That year was 1954.

The Grand Mesa area still remains one of the best spots in the State. Here the snow pack is 108% of average. This will help the water users on the streams originating in that area.

Snow pack on the Uncompahgre Drainage is similar to the Gunnison at 56% of the 1948-62 average.

Mountain soil moisture is better than average for the entire area covered by this report.

The streamflow forecasts issued in this report are revised downward from those issued last month. This is due primarily to the lack of snowfall during April. The Gunnison River is now expected to flow 750,000 acre-feet which is 57% of average. The Uncompahgre River is forecast at 68% of normal for the coming season. Streamflow on the North Fork of the Gunnison should be slightly better and near 75%.

Issued By: Soil Conservation Service

F. A. Mark, State Conservationist,
Colorado

Dearl Beach, Area Conservationist,
Grand Junction, Colorado

"THE CONSERVATION OF WATER BEGINS WITH THE SNOW SURVEY"

| SNOW | | CURRENT INFORMATION | | | PAST RECORD | |
|-------------------|-----|---------------------|---------------------|------------------------|------------------------|------------|
| Snow Course | | Date of Survey | Snow Depth (Inches) | Water Content (Inches) | Water Content (Inches) | |
| | | | | | Last Year | Avg. 48-62 |
| Gunnison River | | | | | | |
| Alexander Lakes | | 4/28 | 62 | 24.4 | 15.7 | 23.0 |
| Black Mesa | | NS | | | - | - |
| Blue Mesa | | 4/27 | 0 | 0.0 | 0 | 2.3* |
| Butte | | 4/27 | 0 | 0.0 | 0 | - |
| Cochetopa Pass | | 4/27 | 0 | 0.0 | 0 | 2.7* |
| Crested Butte | | 4/27 | 0 | 0.0 | 0 | 7.5 |
| Keystone | | 4/27 | 29 | 12.9 | 8.7 | - |
| Lake City | | 4/26 | 0 | 0.0 | 0 | 3.5 |
| Long Gulch | | NS | | | - | - |
| Mesa Lakes | (B) | 4/27 | 36 | 13.7 | 9.6 | 15.9 |
| Monarch Pass | (B) | 4/28 | 14 | 6.8 | 9.3 | 18.4 |
| McClure Pass | | 4/24 | 20 | 7.4 | 2.2 | 10.1* |
| Mineral Creek | (B) | 4/28 | 0 | 0.0 | 11.3 | 12.1* |
| North Lost Trail | (B) | 4/24 | 10 | 3.4 | 1.0 | - |
| Park Cone | | 4/27 | 11 | 4.2 | 5.2 | 8.7 |
| Park Reservoir | | 4/28 | 67 | 27.5 | 21.4 | 25.5 |
| Porphyry Creek | | 4/28 | 22 | 9.9 | 10.9 | 17.7 |
| Tomichi | | 4/28 | 6 | 2.2 | 5.5 | - |
| Trickle Divide | (B) | 4/28 | 68 | 27.7 | 23.8 | 28.8 |
| Uncompahgre River | | | | | | |
| Ironston Park | | 4/27 | 0 | 0.0 | 0 | 7.1 |
| Lizzard Head | | 4/28 | 15 | 6.3 | 13.2 | 13.7 |
| Lone Cone | | 4/27 | 6 | 2.6 | 4.7 | - |
| Red Mountain Pass | (B) | 4/28 | 61 | 25.0 | 25.5 | 31.4 |
| Telluride | | 4/27 | 0 | 0.0 | 0 | 0.7 |
| Trout Lake | | 4/27 | 3 | 1.3 | 3.6 | 9.9* |

RESERVOIR STORAGE (1,000 Acre-Feet)

| Reservoir | Usable Capacity | This Year | Last Year | 15 Year Average 1948-62 |
|-----------|-----------------|-----------|-----------|-------------------------|
| Taylor | 106.2 | 51.8 | 88.7 | 60.3 |

MEASURED FIRST OF MONTH

STREAMFLOW FORECAST (1,000 Acre-Feet)

| Stream and Station | Forecast Period | This Year | Avg. |
|----------------------------|-----------------|-----------|-----------|
| | April - Sept. | % of Avg. | 1948-1962 |
| Gunnison nr Grand Jct. | 750 | 57 | 1305 |
| Surface Creek nr Cedaridge | 15 | 88 | 17 |
| Uncompahgre at Colona | 95 | 68 | 139 |

NOTE: * - 1948-62 (adjusted averages)
 NS - NO SURVEY
 (A) - AIR OBSERVED
 (B) - ON ADJACENT DRAINAGE

SOIL MOISTURE

| Station | Date of Survey | Capacity (Inches) | This Year | Last Year | Avg. All Data |
|---------------|----------------|-------------------|-----------|-----------|---------------|
| Grand Mesa | 4/28 | 12.5 | 9.0 | 12.5 | - |
| King | 4/28 | 3.3 | 3.0 | 3.0 | 2.1 |
| Mineral Creek | 4/28 | 5.7 | 5.4 | 5.6 | 4.1 |
| Placita | 4/28 | 9.3 | 7.8 | 7.5 | 8.1 |

ALL PROFILES 4 FEET DEEP

This Report Prepared by Jack N. Washichek and Donald W. McAndrew, Soil Conservation Service, Colorado State University, Fort Collins, Colo.


RETURN IF NOT DELIVERED
 UNITED STATES
 DEPARTMENT OF AGRICULTURE
 SOIL CONSERVATION SERVICE
 Snow Survey
 Colorado State University
 Fort Collins, Colorado
 OFFICIAL BUSINESS

POSTAGE AND FEES PAID
 U.S. DEPARTMENT OF AGRICULTURE

WATERSHED VII

WATER SUPPLY OUTLOOK
FOR THE SOIL CONSERVATION DISTRICTS IN THE
COLORADO RIVER WATERSHED IN COLORADO
as of
May 1, 1967

U. S. DEPARTMENT OF AGRICULTURE - SOIL CONSERVATION SERVICE
COLORADO EXPERIMENT STATION - STATE ENGINEERS OF COLORADO AND NEW MEXICO



Water users on the Colorado River mainstem should have adequate water this summer. Mountain snow pack now stands at 85% of the 15 year normal. Some of the high elevation snow courses are above normal, due primarily to the late April storms. Two storms dropped as much as three inches of water in some locations. Some melting occurred during the month especially at the lower elevations, however, streamflow has not raised materially. Cool weather could delay runoff for some time.

Mountains soils as of May first contained nearly normal amounts of moisture. Valley soils in the upper basin are in good condition, while the lower areas are reporting only fair soil moisture.

Forecasts on the Colorado mainstem range from 85% of normal at Glenwood to 103% at Granby. This should be adequate for most water users.

The Roaring Fork and Williams Fork Rivers should flow about 85% of the forecast period.

If spring temperatures remain below normal, runoff will be delayed and flows will be less than expected.

Issued By: Soil Conservation Service

F. A. Mark, State Conservationist,
Colorado

D. B. Beach, Area Conservationist,
Grand Junction, Colorado

R. L. Porter, Area Conservationist,
Glenwood Springs, Colorado

"THE CONSERVATION OF WATER BEGINS WITH THE SNOW SURVEY"

SNOW

| Snow Course | Date of Survey | Snow Depth (Inches) | Water Content (Inches) | Water Content (Inches) | |
|---------------------------|----------------|---------------------|------------------------|------------------------|------------|
| | | | | Last Year | Avg. 48-62 |
| <u>Colorado River</u> | | | | | |
| Arrow | 4/26 | 27 | 10.9 | 5.3 | 9.1 |
| Berthoud Pass | 4/25 | 43 | 16.2 | 8.4 | 15.7 |
| Berthoud Summit | 4/27 | 55 | 18.1 | 13.5 | 21.6 |
| Blue River | 4/26 | 4 | 1.1 | 0 | 8.0* |
| Cooper Hill | 4/26 | 43 | 12.6 | 6.8 | - |
| Fiddlers Gulch | 4/28 | 35 | 11.9 | 7.4 | 17.0 |
| Fremont Pass | 4/26 | 57 | 19.5 | 10.3 | 19.5 |
| Frisco | 4/27 | 3 | 1.2 | 1.2 | 5.6* |
| Glen Mar Ranch | 4/24 | 1 | 0.4 | 0.5 | 4.8 |
| Gore Pass | 4/25 | 16 | 5.7 | 1.3 | 7.9* |
| Granby | 4/24 | 10 | 3.7 | 2.3 | 3.3* |
| Grand Lake | 4/26 | 14 | 4.8 | 0.5 | 3.7* |
| Grizzly Peak | 4/27 | 49 | 18.4 | 10.7 | 21.1 |
| Hoosier Pass (B) | 4/26 | 33 | 11.6 | 6.0 | 12.9 |
| Jones Pass | 4/25 | 38 | 15.0 | 9.1 | 16.9* |
| Lake Irene | 4/26 | 61 | 21.8 | 12.0 | 24.7 |
| Lapland | 4/26 | 17 | 6.7 | 0.2 | 9.3 |
| Lulu | 4/28 | 54 | 19.2 | 10.5 | 19.8 |
| Lynx Pass | 4/25 | 22 | 7.4 | 1.6 | 7.8 |
| McKinzie Gulch | 4/26 | 0 | 0.0 | 0 | - |
| Middle Fork Campground | 4/24 | 9 | 3.1 | 4.2 | 6.4 |
| Milner | 4/26 | 38 | 15.1 | 7.4 | 12.1* |
| Monarch Lake | Destroyed | | | 1.3 | 6.4 |
| North Inlet to Grand Lake | 4/26 | 18 | 6.5 | 2.5 | 6.7 |
| Pando | 4/26 | 14 | 6.3 | 3.4 | 8.3 |
| Phantom Valley | 4/26 | 20 | 8.4 | 0.5 | 7.0 |
| Ranch Creek | 4/26 | 28 | 9.4 | 4.2 | 9.6* |
| Shrine Pass | 4/27 | 48 | 18.9 | 9.6 | 20.2 |
| Snake River | 4/28 | 0 | 0.0 | 0 | 5.1* |
| Summit Ranch | 4/25 | 6 | 2.2 | 1.6 | 6.1* |
| Tennessee Pass | 4/28 | 8 | 2.7 | 5.0 | 8.5 |
| Vail Pass | 4/27 | 32 | 14.2 | 6.6 | 16.3* |
| Vasquez Creek | 4/26 | 39 | 12.9 | 8.0 | 14.0 |
| Willow Creek Pass | 4/25 | 35 | 13.5 | 6.5 | 12.0 |
| <u>Roaring Fork River</u> | | | | | |
| Aspen | 4/28 | 45 | 16.1 | 12.0 | - |
| Independence Pass Tunnel | 4/27 | 38 | 17.0 | 10.1 | 17.6 |
| Ivanhoe | 4/27 | 46 | 18.5 | | 19.2 |
| Lift | 4/28 | 53 | 18.2 | 11.4 | 17.8* |
| McClure Pass | 4/24 | 20 | 7.4 | 2.2 | 10.1* |
| Nast | 4/28 | 0 | 0.0 | 0 | 1.7 |
| North Lost Trail | 4/24 | 10 | 3.4 | 1.0 | 8.0 |
| <u>Plateau Creek</u> | | | | | |
| Alexander Lake (B) | 4/28 | 62 | 24.4 | 15.7 | 23.0 |
| Mesa Lakes | 4/27 | 36 | 13.7 | 9.6 | 15.9 |
| Park Reservoir (B) | 4/28 | 67 | 27.5 | 21.4 | 25.5 |
| Trickle Divide | 4/28 | 68 | 27.7 | 23.8 | 28.8 |

SOIL MOISTURE

| Station | Date of Survey | Capacity (Inches) | This Year | Last Year | Avg. All Data |
|----------------|----------------|-------------------|-----------|-----------|---------------|
| Berthoud Pass | 4/25 | 3.9 | 3.2 | 3.5 | 2.8 |
| Blue River | 4/26 | 4.2 | 2.8 | 4.2 | 2.7 |
| Gore | 4/25 | 4.9 | 4.9 | 4.5 | 4.4 |
| Grand Mesa | 4/28 | 12.5 | 9.0 | 12.5 | - |
| Muddy Pass | 4/28 | 11.1 | 9.3 | 11.1 | 8.5 |
| Placita | 4/28 | 9.3 | 7.8 | 7.5 | 8.1 |
| Ranch Creek | 4/26 | 8.7 | 4.9 | 5.9 | 6.5 |
| Vail | 4/27 | 12.3 | 9.0 | 9.1 | 11.0 |
| Vasquez Siphon | 4/26 | 11.0 | 9.4 | 8.6 | 9.2 |

RETURN IF NOT DELIVERED ALL PROFILES 4 FEET DEEP
UNITED STATES

DEPARTMENT OF AGRICULTURE

SOIL CONSERVATION SERVICE

Snow Survey
Colorado State University
Fort Collins, Colorado

OFFICIAL BUSINESS

RESERVOIR STORAGE (1,000 Acre-Feet)

| Reservoir | Usable Capacity | This Year | Last Year | 15 Year Average 1948-62 |
|----------------|-----------------|-----------|-----------|-------------------------|
| Granby | 465.5 | 55.3 | 219.6 | 85.0 |
| Green Mountain | 146.9 | 39.0 | 63.4 | 46.9 |
| Vega | 32.9 | 8.1 | 26.0 | - |
| Williams Fork | 96.8 | 5.9 | 17.9 | - |

MEASURED FIRST OF MONTH

STREAMFLOW FORECAST (1,000 Acre-Feet)

| Stream and Station | Forecast Period April - Sept. | This Year % of Avg. | Avg. 1948-1962 |
|---------------------------------------|-------------------------------|---------------------|----------------|
| Blue River abov. Green Mt. (10) | 220 | 80 | 274 |
| Colo. River nr Granby (11) | 240 | 103 | 233 |
| Colo. River abv Glenwood Springs (12) | 1325 | 85 | 1556 |
| Roaring Fork at Glenwood Springs (14) | 625 | 82 | 762 |
| Williams Fork nr Parshall (15) | 65 | 84 | 77 |
| Willow abv Willow Cr. | 45 | 94 | 48 |
| Colo. nr Cameo (12) | 1950 | 88 | 2213 |

- (10) Observed flow plus change in storage in Dillon Reservoir.
 (11) Observed flow diversions by Adams Tunnel and Grand River Ditch plus change in storage in Granby Reservoir.
 (12) Observed flow plus the changes as indicated in (11) plus Moffat Ditch.
 (14) Observed flow plus diversion through Twin Lakes Tunnel.
 (15) Observed flow plus diversions through Jones Pass Tunnel.

NOTE: * - 1948-62 (adjusted averages)
 NS - NO SURVEY
 (A) - AIR OBSERVED
 (B) - ON ADJACENT DRAINAGE

This Report Prepared by Jack N. Washichek and Donald W. McAndrew, Soil Conservation Service, Soil Conservation Service, Fort Collins, Colo.


POSTAGE AND FEES PAID
U.S. DEPARTMENT OF AGRICULTURE

WATER SUPPLY OUTLOOK
FOR THE SOIL CONSERVATION DISTRICTS IN THE
YAMPA, WHITE, AND NORTH PLATTE
RIVERS WATERSHEDS IN COLORADO

as of
May 1, 1967

WATERSHED VIII

U. S. DEPARTMENT OF AGRICULTURE - SOIL CONSERVATION SERVICE
COLORADO EXPERIMENT STATION - STATE ENGINEERS OF COLORADO AND NEW MEXICO



Water supplies in this area should be adequate this summer. The snow pack on the North Platte, is the best in the state and is 97% of normal.

The Yampa snow pack is only 77% and the White 62%, however, no appreciable water shortage is expected. Snow was still falling over most of this area as of May 1st, so some small increase of the snow pack is expected.

Low elevation snow has started to melt, but the river's flow has not increased yet.

Mountain soils contain about normal moisture, but not as much as last year at this time.

Forecasts assume normal precipitation for the remainder of the season. If this occurs the North Platte should flow 105% of normal, the Yampa and White Rivers about 73%.

The Little Snake should flow just about average.

Issued By: Soil Conservation Service

F. A. Mark, State Conservationist,
Colorado

R. L. Porter, Area Conservationist,
Glenwood Springs, Colorado

"THE CONSERVATION OF WATER BEGINS WITH THE SNOW SURVEY"

| SNOW | | CURRENT INFORMATION | | | PAST RECORD | | STREAMFLOW FORECAST (1,000 Acre-Feet) | | | |
|---------------------------|----------------|---------------------|------------------------|------------------------|-------------|---------------------------|---------------------------------------|---------------------|----------------|--|
| Snow Course | Date of Survey | Snow Depth (Inches) | Water Content (Inches) | Water Content (Inches) | | Stream and Station | Forecast Period April - Sept. | This Year % of Avg. | Avg. 1948-1962 | |
| | | | | Last Year | Avg. 48-62 | | | | | |
| <u>North Platte River</u> | | | | | | | | | | |
| Cameron Pass | 4/26 | 82 | 33.9 | 21.9 | 28.1 | Elk at Clark | 190 | 93 | 205 | |
| Columbine Lodge | 4/28 | 40 | 18.4 | 9.6 | 22.9 | Laramie at Jelm | 121 | 108 | 112 | |
| Deadman Hill (B) | 4/27 | 48 | 17.1 | 13.0 | 18.1 | Little Snake at Lilly | 321 | 100 | 321 | |
| McIntyre (B) | 4/22 | 24 | 9.9 | 5.8 | 10.2* | North Platte at Northgate | 272 | 105 | 260 | |
| Northgate | 4/25 | 8 | 1.9 | 0.4 | 3.0* | White at Meeker | 250 | 75 | 332 | |
| Park View | 4/25 | 19 | 7.1 | 3.0 | 6.8 | Yampa at Maybell | 830 | 90 | 923 | |
| Roach | 4/22 | 51 | 16.8 | 16.4 | 21.0 | Yampa at Steamboat Spr. | 225 | 77 | 292 | |
| Willow Creek Pass (B) | 4/25 | 35 | 13.5 | 6.5 | 12.0 | | | | | |
| <u>Yampa River</u> | | | | | | | | | | |
| Bear River | 4/29 | 16 | 6.0 | 1.1 | 8.3* | | | | | |
| Clark | 4/28 | 7 | 2.8 | 0 | - | | | | | |
| Columbine Lodge (B) | 4/28 | 40 | 18.4 | 9.6 | 22.9 | | | | | |
| Dry Lake | 4/27 | 38 | 14.2 | 6.8 | 17.2 | | | | | |
| Elk River | 4/28 | 38 | 16.3 | 9.3 | 13.4 | | | | | |
| Hahn's Peak | 4/28 | 19 | 8.3 | 1.7 | - | | | | | |
| Lynx Pass | 4/25 | 22 | 7.4 | 1.6 | 7.8 | | | | | |
| Rabbit Ears | 4/28 | 47 | 19.9 | 16.1 | 27.9 | | | | | |
| Yampa View | 4/27 | 0 | 0.0 | 1.3 | 9.7* | | | | | |
| <u>White River</u> | | | | | | | | | | |
| Burro Mountain | 4/25 | 34 | 11.1 | 4.3 | 15.8 | | | | | |
| Rio Blanco | 4/29 | 13 | 5.1 | 0 | 10.5 | | | | | |


| SOIL MOISTURE | | | | | |
|---------------|----------------|-------------------|-----------|-----------|---------------|
| Station | Date of Survey | Capacity (Inches) | This Year | Last Year | Avg. All Data |
| Hahn's Peak | 4/28 | 19.0 | 13.3 | 11.2 | - |
| Laramie Road | 4/30 | 12.4 | 8.1 | 9.1 | 9.0 |
| Muddy Pass | 4/28 | 11.1 | 9.3 | 11.1 | 8.5 |
| Two Mile | 4/27 | 9.1 | 4.3 | 5.5 | 5.6 |
| Willow Pass | 4/25 | 9.5 | 6.5 | 9.5 | 6.9 |

RETURN IF NOT DELIVERED
 UNITED STATES
 DEPARTMENT OF AGRICULTURE
 SOIL CONSERVATION SERVICE
 Snow Survey
 Colorado State University
 Fort Collins, Colorado
 OFFICIAL BUSINESS

POSTAGE AND FEES PAID
 U.S. DEPARTMENT OF AGRICULTURE

WATER SUPPLY OUTLOOK
FOR THE SOIL CONSERVATION DISTRICTS IN THE
LOWER SOUTH PLATTE RIVER WATERSHED IN COLORADO
as of
May 1, 1967

U. S. DEPARTMENT OF AGRICULTURE - SOIL CONSERVATION SERVICE
COLORADO EXPERIMENT STATION - STATE ENGINEERS OF COLORADO AND NEW MEXICO



The snow pack in the mountain watersheds of the South Platte River and its tributaries remains below normal. Even with the big storm of April 13th, the snow pack remains at only 67% of average over the entire basin. There are a few isolated areas in the basin that have a near normal snow pack. These are mostly in the very high elevations.

The water held in the reservoirs throughout the basin remains similar to last month at 107% of average. This water will be an excellent supplement this summer. Following the good rains in the area during this last month most of the irrigated areas are reporting good soil moisture conditions. This situation will help the below normal streamflow as it will lessen the demands on the early flows.

Mountain soil moisture is slightly below normal for this time of year. Some of the snow water will be used to wet up the soil mantle before the spring runoff starts.

Streamflow forecasts range from a high of 90% of average on Clear Creek to a low of 73% on the Cache La Poudre. The Big Thompson, Boulder Creek and Saint Vrain Rivers will flow between 77 to 85% this year. The mainstem of the South Platte will probably flow less than 70% this summer.

Issued By: Soil Conservation Service

F. A. Mark, State Conservationist,
Colorado

J. L. Hall, Area Conservationist,
Sterling, Colorado

"THE CONSERVATION OF WATER BEGINS WITH THE SNOW SURVEY"

SNOW

CURRENT INFORMATION

PAST RECORD

| Snow Course | Date of Survey | Snow Depth (Inches) | Water Content (Inches) | Water Content (Inches) | |
|----------------------------------|----------------------|---------------------------|------------------------------|---------------------------|---------------|
| | | | | Last Year | Avg. 48-62 |
| South Platte River & Tributaries | | | | | |
| Baltimore | 4/27 | 0 | 0 | 0 | - - |
| Berthoud Falls | 4/27 | 19 | 7.2 | 8.4 | 13.8* |
| Big South | 4/29 | 3 | 0.4 | 0 | 0.8 |
| Boulder Falls | 4/29 | 22 | 9.2 | 5.0 | 13.2* |
| Cameron Pass | 4/26 | 82 | 33.9 | 21.9 | 28.1 |
| Chambers Lake | 4/30 | 14 | 6.1 | 0 | 5.5 |
| Como | 4/26 | 6 | 2.1 | - - | - - |
| Copeland Lake | 4/28 | 0 | 0.0 | 0 | 2.3* |
| Deadman Hill | 4/27 | 48 | 17.1 | 13.0 | 18.1 |
| Deer Ridge | 4/27 | 1 | 0.5 | 0 | 3.5* |
| Empire | 4/27 | 22 | 8.4 | 4.2 | 7.1* |
| Geneva Park | 4/26 | 5 | 2.1 | 0.5 | 1.9* |
| Grizzly Peak (B) | 4/27 | 49 | 18.4 | 10.7 | 21.1 |
| Hidden Valley | 4/27 | 29 | 10.6 | 6.9 | 13.6 |
| Hoosier Pass | 4/26 | 33 | 11.6 | 6.0 | 12.9 |
| Horseshoe | 4/26 | 18 | 6.0 | - - | - - |
| Hour Glass Lake | 4/27 | 12 | 4.1 | 1.4 | 7.5 |
| Jefferson Creek | 4/27 | 17 | 5.8 | 1.9 | 8.0* |
| Lake Irene (B) | 4/26 | 61 | 21.8 | 12.0 | 24.7 |
| Long's Peak | 4/29 | 33 | 12.1 | 6.5 | 13.4* |
| Lost Lake | 4/30 | 32 | 7.7 | 1.9 | 10.2* |
| Loveland Lift No. 1 | 4/28 | 76 | 27.5 | 15.1 | - - |
| Loveland Pass | 4/28 | 33 | 13.7 | 4.9 | 16.4 |
| Mosquito | 4/26 | 3 | 1.1 | - - | - - |
| Pine Creek | 4/27 | 0 | 0.0 | 0 | - - |
| Red Feather | | | | 0.7 | 4.9* |
| Two Mile | 4/27 | 53 | 17.3 | 9.8 | 17.8* |
| Trout Creek | 4/28 | 0 | 0.0 | - - | - - |
| University Camp | 4/29 | 35 | 14.1 | 8.2 | 24.9 |
| Ward | 4/27 | 11 | 4.1 | 0.4 | 6.0* |
| Wild Basin | 4/27 | 24 | 7.6 | 5.1 | 14.8 |

RESERVOIR STORAGE (1,000 Acre-Feet)

| Reservoir | Usable Capacity | This Year | Last Year | 15 Year Average 1948-62 |
|----------------|-----------------|-----------|-----------|-------------------------|
| Carter | 108.9 | 95.7 | 107.3 | 79.0 |
| Cheeseman | 79.0 | 31.8 | 77.2 | 54.3 |
| Eleven Mile | 81.9 | 90.9 | 92.3 | 74.6 |
| Empire | 37.7 | 28.3 | 34.1 | 29.6 |
| Horsetooth | 143.5 | 116.8 | 120.3 | 85.6 |
| Jackson | 35.4 | 33.8 | 34.9 | 34.2 |
| Julesburg | 28.2 | 21.7 | 23.2 | 22.0 |
| Point of Rocks | 70.0 | 65.4 | 72.0 | 61.6 |
| Prewitt | 32.8 | 6.3 | 30.2 | 21.7 |
| Riverside | 57.5 | 55.7 | 56.1 | 51.0 |

MEASURED FIRST OF MONTH

STREAMFLOW FORECAST (1,000 Acre-Feet)

| Stream and Station | Forecast Period | This Year | Avg. 1948-1962 |
|------------------------------------|-----------------|-----------|----------------|
| | April - Sept. | % of Avg. | |
| Big Thompson at Drake (2) | 85 | 77 | 110 |
| Boulder at Orodell | 46 | 85 | 54 |
| Cache La Poudre at Canon Mouth (1) | 180 | 73 | 246 |
| Clear Creek at Golden (3) | 120 | 90 | 134 |
| Saint Vrain at Lyons | 65 | 81 | 80 |

SOIL MOISTURE

| Station | Date of Survey | Capacity (Inches) | This Year | Last Year | Avg. All Data |
|---------------|----------------|-------------------|-----------|-----------|---------------|
| Alpine Camp | 4/27 | 6.9 | 3.5 | 4.1 | 4.3 |
| Beaver Dam | 4/27 | 7.3 | 4.8 | 5.2 | 4.7 |
| Clear Creek | 4/28 | 9.5 | 5.8 | 6.4 | 5.9 |
| Feather | 4/28 | 10.1 | 6.5 | 9.4 | 8.1 |
| Guard Station | 4/29 | 6.9 | 4.9 | 4.6 | 4.7 |
| Hoop Creek | 4/26 | 4.9 | 3.5 | 3.5 | 2.9 |
| Hoosier Pass | 4/26 | 7.8 | 4.8 | 6.3 | 5.9 |
| Kenosha Pass | 4/27 | 4.4 | 4.0 | 3.3 | 3.7 |
| Laramie Road | 4/30 | 12.4 | 8.1 | 9.1 | 9.0 |
| Two Mile | 4/27 | 9.1 | 4.3 | 5.5 | 5.6 |

- (1) Observed flow minus diversions from Michigan, Colorado and Laramie Rivers, plus diversions for irrigation and municipal use above station.
- (2) Observed flow plus by-pass to power plants.
- (3) Observed flow minus diversions through Jones Pass.

NOTE: * - 1948-62 (adjusted averages)

NS - NO SURVEY

(A) - AIR OBSERVED

(B) - ON ADJACENT DRAINAGE

This Report Prepared by Jack N. Washichek and Donald W. McAndrew, Soil Conservation Service, Colorado State University, Fort Collins, Colo.

RETURN IF NOT DELIVERED ALL PROFILES 4 FEET DEEP
UNITED STATES

DEPARTMENT OF AGRICULTURE

SOIL CONSERVATION SERVICE

Snow Survey
Colorado State University
Fort Collins, Colorado

OFFICIAL BUSINESS

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF AGRICULTURE

LIST of COOPERATORS

The following organizations cooperate in snow surveys for the Colorado, Platte, Arkansas and Rio Grande watersheds. Many other organizations and individuals furnish valuable information for the snow survey reports. Their cooperation is gratefully acknowledged.

STATE

Colorado State Engineer
New Mexico State Engineer
Nebraska State Engineer
Colorado Experiment Station
Rocky Mountain Forest and Range Experiment Station

FEDERAL

Department of Agriculture

Forest Service
Soil Conservation Service

Department of Interior

Bureau of Reclamation
Geological Survey
National Park Service
Indian Service

Department of Commerce

Weather Bureau

War Department

Army Engineer Corps

Atomic Energy Commission

INVESTOR OWNED UTILITIES

Colorado Public Service Company
Public Service Company of New Mexico

MUNICIPALITIES

| | |
|-----------------|----------------------|
| City of Denver | City of Greeley |
| City of Boulder | City of Fort Collins |

WATER USERS ORGANIZATIONS

Arkansas Valley Ditch Association
Colorado River Water Conservation District

IRRIGATION PROJECTS

Farmers Reservoir and Irrigation Company
San Luis Valley Irrigation District
Santa Maria Reservoir Company
Costilla Land Company
Uncompahgre Valley Water Users' Association
Twin Lakes Reservoir and Canal Company
Trinchera Irrigation Co.

UNITED STATES DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE
SNOW SURVEY UNIT

AG. ENGINEERING SHOP
COLORADO STATE UNIVERSITY
FORT COLLINS, COLORADO 80521

OFFICIAL BUSINESS

U. S. DEPARTMENT OF AGRICULTURE
POSTAGE AND FEES PAID

FIRST CLASS MAIL

FEDERAL - STATE - PRIVATE
COOPERATIVE SNOW SURVEYS

Furnishes the basic data
necessary for forecasting
water supply for irrigation,
domestic and municipal water
supply, hydro-electric power
generation, navigation,
mining and industry

*"The Conservation of Water begins
with the Snow Survey"*